

**HUMAN GLUTATHIONE PEROXIDASE-6**

This application is a **DIVISIONAL** application of U.S. application serial number 09/088,549, filed June 1, 1998, entitled HUMAN GLUTATHIONE PEROXIDASE-6, which is hereby expressly incorporated by reference.

**FIELD OF THE INVENTION**

This invention relates to nucleic acid and amino acid sequences of a human glutathione peroxidase-6 and to the use of these sequences in the diagnosis, treatment, and prevention of reproductive, immune, and cell proliferative disorders.

**BACKGROUND OF THE INVENTION**

Eukaryotic cells extract energy and synthesize macromolecules by a complex series of oxidation-reduction reactions collectively referred to as aerobic metabolism. One consequence of aerobic metabolism is the production of free radicals in the form of superoxides ( $O_2^{\cdot -}$ ) and hydroxyl ions ( $OH^{\cdot}$ ). Superoxides are produced within cells by mitochondria and the endoplasmic reticulum as a consequence of "leakage" of electrons onto  $O_2$  from their correct paths in electron transfer chains. Hydroxyl ions are produced by ionizing radiation and by the reaction of  $O_2^{\cdot -}$  with hydrogen peroxide ( $H_2O_2$ ) at iron- or copper-containing sites. Free radicals, especially hydroxyl ions, are extremely reactive and can interact with almost all molecules, including proteins, carbohydrates, DNA, and lipids. These interactions can lead to the formation of nonradical hydroperoxides, such as phospholipid hydroperoxides. Interaction of hydroxyl ions with DNA may be a significant contributor to the age-dependent development of cancer. Cells also use free radicals and their derivatives in beneficial ways, such as cytochrome P450-mediated oxidations, regulation of smooth muscle tone, and killing of microorganisms by macrophages and granulocytes (Bast, A. et al. (1991) Am. J. Med. 91(3C):2S-13S).

Enzymatic antioxidants act as a defense against free radical damage. Most of the superoxides generated undergo dismutation by superoxide dismutase (SOD) generating hydrogen peroxide.  $H_2O_2$  can also be produced by the action of other oxidase enzymes. Hydrogen peroxide resembles water in its molecular structure and is very diffusible within and between cells. Hydrogen peroxide can be removed by catalase, but the most important enzyme in human cells for removing hydrogen peroxide and hydroperoxides is glutathione peroxidase (GSHPx). GSHPx removes peroxides by using them to oxidize reduced glutathione (GSH) to oxidized

glutathione (GSSG).

The family of glutathione peroxidases encompass three tetrameric glutathione peroxidases (GPx1-3) and the monomeric phospholipid hydroperoxide glutathione peroxidase (PHGPx/GPx4). Although the overall homology between tetrameric enzymes and GPx4 is less than 30%, a pronounced similarity has been detected in clusters involved in the active site and a common catalytic triad has been defined by structural and kinetic data (Epp, O. et al. (1983) Eur. J. Biochem. 133:51-69). The family members show different tissue distributions. GPx1 is ubiquitously expressed in cells, whereas GPx2 is present in the liver and colon and GPx3 in blood plasma. GPx4 is found at low level in all tissues but is expressed at high level in the testis. These tissue localization patterns may be important for regulating the level and targets of glutathione peroxidase activity (Ursini, F. et al (1995) Methods Enzymol. 252:38-53).

GPx4 is unique in both its structure and activity. GPx4 is the only monomeric glutathione peroxidase found in mammals. It is also the only mammalian glutathione peroxidase to show high affinity for and reactivity with phospholipid hydroperoxides, and to be membrane associated. The inhibition of lipid peroxidation by GPx4 requires glutathione and physiological levels of vitamin E, suggesting a tandem mechanism for the antioxidant activities of GPx4 and vitamin E. GPx4 also has alternative transcription and translation start sites which determine its subcellular localization (Esworthy, R.S. et al. (1994) Gene 144:317-318; and Maiorino, M. et al. (1990) Methods Enzymol. 186:448-450).

Antioxidant defenses exist as a balanced, coordinated system. Disruption in this balance leads to clinical manifestations. For example, an excess of SOD in relation to GSHPx may be relevant to the clinical condition known as Down's Syndrome. Low antioxidant levels or high  $O_2^-$  and  $H_2O_2$  levels produce oxidative stress. Oxidative stress induced by phagocytes at sites of chronic inflammation lead to rheumatoid arthritis in the joints and inflammatory bowel diseases in the intestine. Asthma is also a manifestation of an inflammatory reaction in the lung and is related to oxygen free radical formation (Sies, H. (1991) Am. J. Med. 91 (3C):31S-38S).

The discovery of a new glutathione peroxidase and the polynucleotides encoding it satisfies a need in the art by providing new compositions which are useful in the diagnosis, treatment, and prevention of reproductive, immune, and cell proliferative disorders.

## SUMMARY OF THE INVENTION

The invention is based on the discovery of a new human glutathione peroxidase (GPx6), the polynucleotides encoding GPx6, and the use of these compositions for the diagnosis,

treatment, or prevention of reproductive, immune, and cell proliferative disorders.

The invention features a substantially purified polypeptide comprising the amino acid sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1.

5 The invention further provides a substantially purified variant having at least 90% amino acid sequence identity to the amino acid sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1. The invention also provides an isolated and purified polynucleotide encoding the polypeptide comprising the sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1. The invention also includes an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide encoding the polypeptide comprising the  
10 amino acid sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1.

The invention further provides an isolated and purified polynucleotide which hybridizes under stringent conditions to the polynucleotide encoding the polypeptide comprising the amino acid sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1, as well as an isolated and purified polynucleotide which is complementary to the polynucleotide encoding the polypeptide  
15 comprising the amino acid sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1.

The invention also provides an isolated and purified polynucleotide comprising the polynucleotide sequence of SEQ ID NO:2 or a fragment of SEQ ID NO:2, and an isolated and purified polynucleotide variant having at least 90% polynucleotide sequence identity to the polynucleotide comprising the polynucleotide sequence of SEQ ID NO:2 or a fragment of SEQ  
20 ID NO:2. The invention also provides an isolated and purified polynucleotide having a sequence complementary to the polynucleotide comprising the polynucleotide sequence of SEQ ID NO:2 or a fragment of SEQ ID NO:2.

The invention further provides an expression vector comprising at least a fragment of the polynucleotide encoding the polypeptide comprising the sequence of SEQ ID NO:1 or a fragment  
25 of SEQ ID NO:1. In another aspect, the expression vector is contained within a host cell.

The invention also provides a method for producing a polypeptide comprising the amino acid sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1, the method comprising the steps of: (a) culturing the host cell comprising an expression vector containing at least a fragment of a polynucleotide encoding the polypeptide comprising the amino acid sequence of SEQ ID NO:1  
30 or a fragment of SEQ ID NO: 1 under conditions suitable for the expression of the polypeptide; and (b) recovering the polypeptide from the host cell culture.

The invention also provides a pharmaceutical composition comprising a substantially purified polypeptide having the sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1 in conjunction with a suitable pharmaceutical carrier.

The invention further includes a purified antibody which binds to a polypeptide comprising the sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1, as well as a purified agonist and a purified antagonist of the polypeptide.

The invention also provides a method for treating or preventing a reproductive disorder, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising substantially purified polypeptide having the amino acid sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1.

The invention also provides a method for treating or preventing an immune disorder, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising substantially purified polypeptide having the amino acid sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1.

The invention also provides a method for treating or preventing a cell proliferative disorder, the method comprising administering to a subject in need of such treatment an effective amount of a pharmaceutical composition comprising substantially purified polypeptide having the amino acid sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1.

The invention also provides a method for detecting a polynucleotide encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 1 or a fragment of SEQ ID NO: 1 in a biological sample containing nucleic acids, the method comprising the steps of: (a) hybridizing the complement of the polynucleotide encoding the polypeptide comprising the amino acid sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1 to at least one of the nucleic acids of the biological sample, thereby forming a hybridization complex; and (b) detecting the hybridization complex, wherein the presence of the hybridization complex correlates with the presence of a polynucleotide encoding the polypeptide comprising the amino acid sequence of SEQ ID NO:1 or a fragment of SEQ ID NO:1 in the biological sample. In one aspect, the nucleic acids of the biological sample are amplified by the polymerase chain reaction prior to the hybridizing step.

## BRIEF DESCRIPTION OF THE FIGURES

Figures 1A, 1B, and 1C show the amino acid sequence (SEQ ID NO:1) and nucleic acid sequence (SEQ ID NO:2) of GPx6. The alignment was produced using MacDNASIS PRO software (Hitachi Software Engineering Co. Ltd., San Bruno, CA).

Figure 2 shows the amino acid sequence alignments among GPx6 (Incyte Clone number 1817518; SEQ ID NO:1), and Human phospholipid hydroperoxide glutathione peroxidase (GI 544432; SEQ ID NO:3), produced using the multisequence alignment program of LASERGENE

software (DNASTAR Inc, Madison WI).

## DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular methodology, protocols, cell lines, vectors, and reagents described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, devices, and materials are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, vectors, and methodologies which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

## DEFINITIONS

"GPx6," as used herein, refers to the amino acid sequences of substantially purified GPx6 obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and preferably the human species, from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist," as used herein, refers to a molecule which, when bound to GPx6, increases or prolongs the duration of the effect of GPx6. Agonists may include proteins, nucleic acids, carbohydrates, or any other molecules which bind to and modulate the effect of GPx6.

An "allelic variant," as this term is used herein, is an alternative form of the gene encoding GPx6. Allelic variants may result from at least one mutation in the nucleic acid

sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding GPx6, as described herein, include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polynucleotide the same as GPx6 or a polypeptide with at least one functional characteristic of GPx6. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding GPx6, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding GPx6. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent GPx6. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of GPx6 is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, positively charged amino acids may include lysine and arginine, and amino acids with uncharged polar head groups having similar hydrophilicity values may include leucine, isoleucine, and valine; glycine and alanine; asparagine and glutamine; serine and threonine; and phenylalanine and tyrosine.

The terms "amino acid" or "amino acid sequence," as used herein, refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. In this context, "fragments," "immunogenic fragments," or "antigenic fragments" refer to fragments of GPx6 which are preferably about 5 to about 15 amino acids in length, most preferably 14 amino acids, and which retain some biological activity or immunological activity of GPx6. Where "amino acid sequence" is recited herein to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification," as used herein, relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art. (See, e.g., Dieffenbach, C.W. and G.S. Dveksler (1995) PCR

Primer, a Laboratory Manual, Cold Spring Harbor Press, Plainview, NY, pp.1-5.)

The term "antagonist," as it is used herein, refers to a molecule which, when bound to GPx6, decreases the amount or the duration of the effect of the biological or immunological activity of GPx6. Antagonists may include proteins, nucleic acids, carbohydrates, antibodies, or any other molecules which decrease the effect of GPx6.

As used herein, the term "antibody" refers to intact molecules as well as to fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv fragments, which are capable of binding the epitopic determinant. Antibodies that bind GPx6 polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant," as used herein, refers to that fragment of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (given regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense," as used herein, refers to any composition containing a nucleic acid sequence which is complementary to the "sense" strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and to block either transcription or translation. The designation "negative" can refer to the antisense strand, and the designation "positive" can refer to the sense strand.

As used herein, the term "biologically active," refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic GPx6, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" or "complementarity," as used herein, refer to the natural binding of polynucleotides by base pairing. For example, the sequence "A-G-T" binds to the complementary sequence "T-C-A." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acids strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" or a "composition comprising a given amino acid sequence," as these terms are used herein, refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation, an aqueous solution, or a sterile composition. Compositions comprising polynucleotide sequences encoding GPx6 or fragments of GPx6 may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts, e.g., NaCl, detergents, e.g., sodium dodecyl sulfate (SDS), and other components, e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.

"Consensus sequence," as used herein, refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using XL-PCR (Perkin Elmer, Norwalk, CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of more than one Incyte Clone using a computer program for fragment assembly, such as the GELVIEW Fragment Assembly system (GCG, Madison, WI). Some sequences have been both extended and assembled to produce the consensus sequence.

As used herein, the term "correlates with expression of a polynucleotide" indicates that the detection of the presence of nucleic acids, the same or related to a nucleic acid sequence encoding GPx6, by Northern analysis is indicative of the presence of nucleic acids encoding GPx6 in a sample, and thereby correlates with expression of the transcript from the polynucleotide encoding GPx6.

A "deletion," as the term is used herein, refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative," as used herein, refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence



can include, for example, replacement of hydrogen by an alkyl, acyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

The term "similarity," as used herein, refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or Northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" or "% identity" refer to the percentage of sequence similarity found in a comparison of two or more amino acid or nucleic acid sequences. Percent identity can be determined electronically, e.g., by using the MEGALIGN program (DNASTAR, Inc., Madison WI). The MEGALIGN program can create alignments between two or more sequences according to different methods, e.g., the clustal method. (See, e.g., Higgins, D.G. and P.M. Sharp (1988) *Gene* 73:237-244.) The clustal algorithm groups sequences into clusters by examining the distances between all pairs. The clusters are aligned pairwise and then in groups. The percentage similarity between two amino acid sequences, e.g., sequence A and sequence B, is calculated by dividing the length of sequence A, minus the number of gap residues in sequence A, minus the number of gap residues in sequence B, into the sum of the residue matches between sequence A and sequence B, times one hundred. Gaps of low or of no similarity between the two amino acid sequences are not included in determining percentage similarity. Percent identity between nucleic acid sequences can also be counted or calculated by other methods known in the

art, e.g., the Jotun Hein method. (See, e.g., Hein, J. (1990) *Methods Enzymol.* 183:626-645.) Identity between sequences can also be determined by other methods known in the art, e.g., by varying hybridization conditions.

“Human artificial chromosomes” (HACs), as described herein, are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance. (See, e.g., Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.)

The term “humanized antibody,” as used herein, refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

“Hybridization,” as the term is used herein, refers to any process by which a strand of nucleic acid binds with a complementary strand through base pairing.

As used herein, the term “hybridization complex” refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g.,  $C_0t$  or  $R_0t$  analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words “insertion” or “addition,” as used herein, refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively, to the sequence found in the naturally occurring molecule.

“Immune response” can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

The term “microarray,” as used herein, refers to an arrangement of distinct polynucleotides arrayed on a substrate, e.g., paper, nylon or any other type of membrane, filter, chip, glass slide, or any other suitable solid support.

The terms “element” or “array element” as used herein in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

The term “modulate,” as it appears herein, refers to a change in the activity of GPx6. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of GPx6.

The phrases “nucleic acid” or “nucleic acid sequence,” as used herein, refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material. In this context, “fragments” refers to those nucleic acid sequences which, when translated, would produce polypeptides retaining some functional characteristic, e.g., antigenicity, or structural domain characteristic, e.g., ATP-binding site, of the full-length polypeptide.

The terms “operably associated” or “operably linked,” as used herein, refer to functionally related nucleic acid sequences. A promoter is operably associated or operably linked with a coding sequence if the promoter controls the translation of the encoded polypeptide. While operably associated or operably linked nucleic acid sequences can be contiguous and in the same reading frame, certain genetic elements, e.g., repressor genes, are not contiguously linked to the sequence encoding the polypeptide but still bind to operator sequences that control expression of the polypeptide.

The term “oligonucleotide,” as used herein, refers to a nucleic acid sequence of at least about 6 nucleotides to 60 nucleotides, preferably about 15 to 30 nucleotides, and most preferably about 20 to 25 nucleotides, which can be used in PCR amplification or in a hybridization assay or microarray. As used herein, the term “oligonucleotide” is substantially equivalent to the terms “amplimer,” “primer,” “oligomer,” and “probe,” as these terms are commonly defined in the art.

“Peptide nucleic acid” (PNA), as used herein, refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell. (See, e.g., Nielsen, P.E. et al. (1993) *Anticancer Drug Des.* 8:53-63.)

The term “sample,” as used herein, is used in its broadest sense. A biological sample suspected of containing nucleic acids encoding GPx6, or fragments thereof, or GPx6 itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a solid support; a tissue; a tissue print; etc.

As used herein, the terms “specific binding” or “specifically binding” refer to that interaction between a protein or peptide and an agonist, an antibody, or an antagonist. The

interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will  
 5 reduce the amount of labeled A that binds to the antibody.

As used herein, the term "stringent conditions" refers to conditions which permit hybridization between polynucleotides and the claimed polynucleotides. Stringent conditions can be defined by salt concentration, the concentration of organic solvent (e.g., formamide), temperature, and other conditions well known in the art. In particular, stringency can be  
 10 increased by reducing the concentration of salt, increasing the concentration of formamide, or raising the hybridization temperature.

The term "substantially purified," as used herein, refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least about 60% free, preferably about 75% free, and most preferably about 90% free from  
 15 other components with which they are naturally associated.

A "substitution," as used herein, refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Transformation," as defined herein, describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which  
 20 inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "variant" of GPx6, as used herein, refers to an amino acid sequence that is altered by one or more amino acids. The variant may have "conservative" changes, wherein a substituted  
 30 amino acid has similar structural or chemical properties (e.g., replacement of leucine with isoleucine). More rarely, a variant may have "nonconservative" changes (e.g., replacement of glycine with tryptophan). Analogous minor variations may also include amino acid deletions or insertions, or both. Guidance in determining which amino acid residues may be substituted,

inserted, or deleted without abolishing biological or immunological activity may be found using computer programs well known in the art, for example, LASERGENE software.

## THE INVENTION

The invention is based on the discovery of a new human glutathione peroxidase (GPx6), the polynucleotides encoding GPx6, and the use of these compositions for the diagnosis, treatment, or prevention of reproductive, immune, and cell proliferative disorders.

Nucleic acids encoding the GPx6 of the present invention were first identified in Incyte Clone 1817518 from the prostate cDNA library (PROSNOT20) using a computer search, e.g., BLAST, for amino acid sequence alignments. A consensus sequence, SEQ ID NO:2, was derived from the following overlapping and/or extended nucleic acid sequences: Incyte Clones 3321614H1 (PTHYNOT03), 1817518H1 and 1817518CT1 (PROSNOT20), and 366321H1 (SYNORAT01).

In one embodiment, the invention encompasses a polypeptide comprising the amino acid sequence of SEQ ID NO:1, as shown in Figures 1A, 1B, and 1C. GPx6 is 187 amino acids in length and has three potential casein kinase II phosphorylation sites at residues S96, S136, and S164; two potential protein kinase C phosphorylation sites at S96 and S136; and a glutathione peroxidase signature at L82. PFAM identifies significant sequence identity with glutathione peroxidase. BLOCKS identifies significant sequence identity with glutathione peroxidase over residues 35-60, 63-100, 107-133, and 138-159, which includes the glutathione peroxidase catalytic triad formed by residues Cys57, Gln92, and Trp142. As shown in Figure 2, GPx6 has chemical and structural similarity with human phospholipid hydroperoxide glutathione peroxidase (GPx4) (SEQ ID NO:3). In particular, GPx6 and GPx4 share 40% identity, and have similar molecular weights (20.9 kD and 19.5 kD, respectively) and isoelectric points (7.79 and 7.88, respectively). Northern analysis shows the expression of this sequence in various libraries, at least 38% of which are immortalized or cancerous, at least 33% of which are associated with fetal development and cell proliferation, and at least 14% of which involve immune response. Of particular note is the expression of GPx6 in nervous, reproductive, and developmental tissues.

The invention also encompasses GPx6 variants. A preferred GPx6 variant is one which has at least about 80%, more preferably at least about 90%, and most preferably at least about 95% amino acid sequence identity to the GPx6 amino acid sequence, and which contains at least one functional or structural characteristic of GPx6.

The invention also encompasses polynucleotides which encode GPx6. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising the sequence of

SEQ ID NO:2, which encodes an GPx6.

The invention also encompasses a variant of a polynucleotide sequence encoding Gpx6. In particular, such a variant polynucleotide sequence will have at least about 80%, more preferably at least about 90%, and most preferably at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding GPx6. A particular aspect of the invention encompasses a variant of SEQ ID NO:2 which has at least about 80%, more preferably at least about 90%, and most preferably at least about 95% polynucleotide sequence identity to SEQ ID NO:2. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of GPx6.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding GPx6, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring GPx6, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode GPx6 and its variants are preferably capable of hybridizing to the nucleotide sequence of the naturally occurring GPx6 under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding GPx6 or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding GPx6 and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode GPx6 and GPx6 derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding GPx6 or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:2, or a fragment of SEQ ID NO:2, under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987)

5 *Methods Enzymol.* 152:507-511.) For example, stringent salt concentration will ordinarily be less than about 750 mM NaCl and 75 mM trisodium citrate, preferably less than about 500 mM NaCl and 50 mM trisodium citrate, and most preferably less than about 250 mM NaCl and 25 mM trisodium citrate. Low stringency hybridization can be obtained in the absence of organic solvent, e.g., formamide, while high stringency hybridization can be obtained in the presence of  
 10 at least about 35% formamide, and most preferably at least about 50% formamide. Stringent temperature conditions will ordinarily include temperatures of at least about 30°C, more preferably of at least about 37°C, and most preferably of at least about 42°C. Varying additional parameters, such as hybridization time, the concentration of detergent, e.g., sodium dodecyl sulfate (SDS), and the inclusion or exclusion of carrier DNA, are well known to those skilled in  
 15 the art. Various levels of stringency are accomplished by combining these various conditions as needed. In a preferred embodiment, hybridization will occur at 30°C in 750 mM NaCl, 75 mM trisodium citrate, and 1% SDS. In a more preferred embodiment, hybridization will occur at 37°C in 500 mM NaCl, 50 mM trisodium citrate, 1% SDS, 35% formamide, and 100 µg/ml denatured salmon sperm DNA (ssDNA). In a most preferred embodiment, hybridization will  
 20 occur at 42°C in 250 mM NaCl, 25 mM trisodium citrate, 1% SDS, 50 % formamide, and 200 µg/ml ssDNA. Useful variations on these conditions will be readily apparent to those skilled in the art.

The washing steps which follow hybridization can also vary in stringency. Wash stringency conditions can be defined by salt concentration and by temperature. As above, wash  
 25 stringency can be increased by decreasing salt concentration or by increasing temperature. For example, stringent salt concentration for the wash steps will preferably be less than about 30 mM NaCl and 3 mM trisodium citrate, and most preferably less than about 15 mM NaCl and 1.5 mM trisodium citrate. Stringent temperature conditions for the wash steps will ordinarily include temperature of at least about 25°C, more preferably of at least about 42°C, and most preferably of  
 30 at least about 68°C. In a preferred embodiment, wash steps will occur at 25°C in 30 mM NaCl, 3 mM trisodium citrate, and 0.1% SDS. In a more preferred embodiment, wash steps will occur at 42°C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1% SDS. In a most preferred embodiment, wash steps will occur at 68°C in 15 mM NaCl, 1.5 mM trisodium citrate, and 0.1%

SDS. Additional variations on these conditions will be readily apparent to those skilled in the art.

Methods for DNA sequencing are well known and generally available in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, Sequenase® (US Biochemical Corp.,

- 5 Cleveland, OH), Taq polymerase (Perkin Elmer), thermostable T7 polymerase (Amersham, Chicago, IL), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE Amplification System (GIBCO BRL, Gaithersburg, MD). Preferably, the process is automated with machines such as the Hamilton Micro Lab 2200 (Hamilton, Reno, NV), Peltier Thermal Cycler (PTC200; MJ Research, Watertown, MA) and the ABI Catalyst and
- 10 373 and 377 DNA Sequencers (Perkin Elmer).

- The nucleic acid sequences encoding GPx6 may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence
- 15 from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) PCR Methods Applic. 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) Nucleic Acids Res. 16:8186.) A third method, capture PCR, involves
  - 20 PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al. (1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See,
  - 25 e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060). Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries to walk genomic DNA (Clontech, Palo Alto, CA). This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth,
  - 30 MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been



size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode GPx6 may be cloned in recombinant DNA molecules that direct expression of GPx6, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express GPx6.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter GPx6-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In another embodiment, sequences encoding GPx6 may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223, and Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.) Alternatively, GPx6 itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques.

(See, e.g., Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A Peptide Synthesizer (Perkin Elmer). Additionally, the amino acid sequence of GPx6, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

5 The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman and Co., New York, NY.)

10 In order to express a biologically active GPx6, the nucleotide sequences encoding GPx6 or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in  
15 polynucleotide sequences encoding GPx6. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding GPx6. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding GPx6 and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no  
20 additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the  
25 particular host cell system used. (See, e.g., Scharf, D. et al. (1994) Results Probl. Cell Differ. 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding GPx6 and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques,  
30 synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview, NY, ch. 4, 8, and 16-17; and Ausubel, F.M. et al. (1995, and periodic supplements) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding GPx6. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus (CaMV) or tobacco mosaic virus (TMV)) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding GPx6. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding GPx6 can be achieved using a multifunctional E. coli vector such as BLUESCRIPT (Stratagene) or PSPORT1 plasmid (GIBCO BRL). Ligation of sequences encoding GPx6 into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of GPx6 are needed, e.g. for the production of antibodies, vectors which direct high level expression of GPx6 may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of GPx6. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, supra; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; Scorer, C.A. et al. (1994) Bio/Technology 12:181-184.)

Plant systems may also be used for expression of GPx6. Transcription of sequences encoding GPx6 may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV. (Takamatsu, N. (1987) EMBO J. 6:307-311.) Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al. (1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell

Differ. 17:85-105.) These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., Hobbs, S. or Murry, L.E. in McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York, NY; pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding GPx6 may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses GPx6 in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes.

For long term production of recombinant proteins in mammalian systems, stable expression of GPx6 in cell lines is preferred. For example, sequences encoding GPx6 can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *apv* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; and Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* or *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570;

Colbere-Garapin, F. et al (1981) *J. Mol. Biol.* 150:1-14; and Murry, *supra.*) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) *Proc. Natl. Acad. Sci. USA* 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP) (Clontech, Palo Alto, CA),  $\beta$  glucuronidase and its substrate  $\beta$ -D-glucuronoside, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. et al. (1995) *Methods Mol. Biol.* 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding GPx6 is inserted within a marker gene sequence, transformed cells containing sequences encoding GPx6 can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding GPx6 under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding GPx6 and that express GPx6 may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of GPx6 using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on GPx6 is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) *Serological Methods, a Laboratory Manual*, APS Press, St. Paul, MN, Section IV; Coligan, J.E. et al. (1997 and periodic supplements) *Current Protocols in Immunology*, Greene Pub. Associates and Wiley-Interscience, New York, NY; and Maddox, D.E. et al. (1983) *J. Exp. Med.* 158:1211-1216).

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled

hybridization or PCR probes for detecting sequences related to polynucleotides encoding GPx6 include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding GPx6, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Pharmacia & Upjohn (Kalamazoo, MI), Promega (Madison, WI), and U.S. Biochemical Corp. (Cleveland, OH). Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding GPx6 may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode GPx6 may be designed to contain signal sequences which direct secretion of GPx6 through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding GPx6 may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric GPx6 protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of GPx6 activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to,

glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the GPx6 encoding sequence and the heterologous protein sequence, so that GPx6 may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel, F.M. et al. (1995 and periodic supplements) Current Protocols in Molecular Biology, John Wiley & Sons, New York, NY, ch 10. A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled GPx6 may be achieved in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract systems (Promega, Madison, WI). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, preferably <sup>35</sup>S-methionine.

Fragments of GPx6 may be produced not only by recombinant production, but also by direct peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Various fragments of GPx6 may be synthesized separately and then combined to produce the full length molecule.

## THERAPEUTICS

Chemical and structural similarity exists between GPx6 and phospholipid hydroperoxide glutathione peroxidase from human (GI544432). In addition, GPx6 is expressed in reproductive and inflamed tissues, and in proliferative tissues, including those associated with development and cancer. Therefore, GPx6 appears to play a role in reproductive and immune disorders, and cell proliferative disorders including developmental disorders and cancer.

Therefore, in one embodiment, GPx6 or a fragment or derivative thereof may be administered to a subject to treat or prevent a reproductive disorder. Such reproductive disorders can include, but are not limited to, disorders of prolactin production; infertility, including tubal

disease, ovulatory defects, and endometriosis; disruptions of the estrous cycle, disruptions of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, endometrial and ovarian tumors, uterine fibroids, autoimmune disorders, ectopic pregnancies, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; disruptions of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, carcinoma of the male breast, and gynecomastia.

In another embodiment, a vector capable of expressing GPx6 or a fragment or derivative thereof may be administered to a subject to treat or prevent a reproductive disorder including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified GPx6 in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a reproductive disorder including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of GPx6 may be administered to a subject to treat or prevent a reproductive disorder including, but not limited to, those listed above.

In another embodiment, GPx6 or a fragment or derivative thereof may be administered to a subject to treat or prevent an immune disorder. Such immune disorders can include, but are not limited to, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma.

In another embodiment, a vector capable of expressing GPx6 or a fragment or derivative thereof may be administered to a subject to treat or prevent an immune disorder including, but



not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified GPx6 in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent an immune disorder including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of GPx6 may be administered to a subject to treat or prevent an immune disorder including, but not limited to, those listed above.

In another embodiment, GPx6 or a fragment or derivative thereof may be administered to a subject to treat or prevent a cell proliferative or developmental disorder. Such cell proliferative or developmental disorders can include, but are not limited to, actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; and developmental disorders which can include, but are not limited to, renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucocutaneous dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, and sensorineural hearing loss.

In another embodiment, a vector capable of expressing GPx6 or a fragment or derivative thereof may be administered to a subject to treat or prevent a cell proliferative or developmental disorder including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified GPx6 in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a cell proliferative or developmental disorder including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of GPx6 may be

administered to a subject to treat or prevent a cell proliferative or developmental disorder including, but not limited to, those listed above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of GPx6 may be produced using methods which are generally known in the art. In particular, purified GPx6 may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind GPx6. Antibodies to GPx6 may also be generated using methods that are well known in the art. Such antibodies may include, but are not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are especially preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with GPx6 or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to GPx6 have an amino acid sequence consisting of at least about 5 amino acids, and, more preferably, of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of GPx6 amino acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to GPx6 may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are

not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique. (See, e.g., Kohler, G. et al. (1975) *Nature* 256:495-497; Kozbor, D. et al. (1985) *J. Immunol. Methods* 81:31-42; Cote, R.J. et al. (1983) *Proc. Natl. Acad. Sci. USA* 80:2026-2030; and Cole, S.P. et al. (1984) *Mol. Cell Biol.* 62:109-120.)

- 5 In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6851-6855; Neuberger, M.S. et al. (1984) *Nature* 312:604-608; and Takeda, S. et al. (1985) *Nature* 314:452-454.) Alternatively, techniques described for the
- 10 production of single chain antibodies may be adapted, using methods known in the art, to produce GPx6-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton D.R. (1991) *Proc. Natl. Acad. Sci. USA* 88:10134-10137.)

- 15 Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86:3833-3837; and Winter, G. et al. (1991) *Nature* 349:293-299.)

- Antibody fragments which contain specific binding sites for GPx6 may also be generated.
- 20 For example, such fragments include, but are not limited to, F(ab')<sub>2</sub> fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')<sub>2</sub> fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) *Science* 246:1275-1281.)

- 25 Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between GPx6 and its specific antibody. A two-site, monoclonal-based immunoassay utilizing
- 30 monoclonal antibodies reactive to two non-interfering GPx6 epitopes is preferred, but a competitive binding assay may also be employed. (Maddox, supra.)

In another embodiment of the invention, the polynucleotides encoding GPx6, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the

complement of the polynucleotide encoding GPx6 may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding GPx6. Thus, complementary molecules or fragments may be used to modulate GPx6 activity, or to achieve regulation of gene function.

- 5 Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding GPx6.

Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding GPx6. (See, e.g., Sambrook, supra; and Ausubel, supra.)

Genes encoding GPx6 can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding GPx6. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding GPx6. Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing Co., Mt. Kisco, NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the

ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding GPx6.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding GPx6. Such DNA sequences may be incorporated into a wide variety of vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends of the molecule, or the use of phosphorothioate or 2'-O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of

such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

An additional embodiment of the invention relates to the administration of a pharmaceutical or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of GPx6, antibodies to GPx6, and mimetics, agonists, antagonists, or inhibitors of GPx6. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired,

disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol.

Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acid. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2%

sucrose, and 2% to 7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of GPx6, such labeling would include amount, frequency, and method of administration.

Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example GPx6 or fragments thereof, antibodies of GPx6, and agonists, antagonists or inhibitors of GPx6, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the  $ED_{50}$  (the dose therapeutically effective in 50% of the population) or  $LD_{50}$  (the dose lethal to 50% of the population) statistics. The dose ratio of therapeutic to toxic effects is the therapeutic index, and it can be expressed as the  $ED_{50}/LD_{50}$  ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the  $ED_{50}$  with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance



rate of the particular formulation.

Normal dosage amounts may vary from about 0.1  $\mu\text{g}$  to 100,000  $\mu\text{g}$ , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art.

- 5 Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

## DIAGNOSTICS

- 10 In another embodiment, antibodies which specifically bind GPx6 may be used for the diagnosis of disorders characterized by expression of GPx6, or in assays to monitor patients being treated with GPx6 or agonists, antagonists, or inhibitors of GPx6. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for GPx6 include methods which utilize the antibody and a label to detect
- 15 GPx6 in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

- A variety of protocols for measuring GPx6, including ELISAs, RIAs, and FACS, are
- 20 known in the art and provide a basis for diagnosing altered or abnormal levels of GPx6 expression. Normal or standard values for GPx6 expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, preferably human, with antibody to GPx6 under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, preferably by photometric means. Quantities
- 25 of GPx6 expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

- In another embodiment of the invention, the polynucleotides encoding GPx6 may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide
- 30 sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantitate gene expression in biopsied tissues in which expression of GPx6 may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of GPx6, and to monitor regulation of GPx6 levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding GPx6 or closely related molecules may be used to identify nucleic acid sequences which encode GPx6. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification (maximal, high, intermediate, or low), will determine whether the probe identifies only naturally occurring sequences encoding GPx6, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and should preferably have at least 50% sequence identity to any of the GPx6 encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:2 or from genomic sequences including promoters, enhancers, and introns of the GPx6 gene.

Means for producing specific hybridization probes for DNAs encoding GPx6 include the cloning of polynucleotide sequences encoding GPx6 or GPx6 derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as  $^{32}\text{P}$  or  $^{35}\text{S}$ , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding GPx6 may be used for the diagnosis of a disorder associated with expression of GPx6. Examples of such a disorder include, but are not limited to, reproductive disorders such as disorders of prolactin production; infertility, including tubal disease, ovulatory defects, and endometriosis; disruptions of the estrous cycle, disruptions of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, endometrial and ovarian tumors, uterine fibroids, autoimmune disorders, ectopic pregnancies, and teratogenesis; cancer of the breast, fibrocystic breast disease, and galactorrhea; disruptions of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, carcinoma of the male breast, gynecomastia; immune disorders such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis,

dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation,

5 osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; and cell proliferative disorders such

10 as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers

15 of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus; and developmental disorders such as renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis

20 syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, and sensorineural hearing loss. The polynucleotide sequences encoding GPx6 may be

25 used in Southern or Northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and ELISA assays; and in microarrays utilizing fluids or tissues from patients to detect altered GPx6 expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding GPx6 may be useful in assays

30 that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding GPx6 may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantitated and compared with a standard value. If the amount of signal in the patient sample is significantly

altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding GPx6 in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

5 In order to provide a basis for the diagnosis of a disorder associated with expression of GPx6, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding GPx6, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from  
10 normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

15 Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

20 With respect to cancer, the presence of a relatively high amount of transcript in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

25 Additional diagnostic uses for oligonucleotides designed from the sequences encoding GPx6 may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding GPx6, or a fragment of a polynucleotide complementary to the polynucleotide encoding GPx6, and will be employed under optimized conditions for  
30 identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantitation of closely related DNA or RNA sequences.

Methods which may also be used to quantitate the expression of GPx6 include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and

interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) *J. Immunol. Methods* 159:235-244; and Duplaa, C. et al. (1993) *Anal. Biochem.* 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in an ELISA format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) *Proc. Natl. Acad. Sci. USA* 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) *Proc. Natl. Acad. Sci. USA* 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

In another embodiment of the invention, nucleic acid sequences encoding GPx6 may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Price, C.M. (1993) *Blood Rev.* 7:127-134; and Trask, B.J. (1991) *Trends Genet.* 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, R.A. (ed.) Molecular Biology and Biotechnology, VCH Publishers New York, NY, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) site. Correlation between the location of the gene encoding GPx6 on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

In situ hybridization of chromosomal preparations and physical mapping techniques, such

as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) *Nature* 336:577-580.) The nucleotide sequence of the subject invention may also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, GPx6, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between GPx6 and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. The test compounds are reacted with GPx6, or fragments thereof, and washed. Bound GPx6 is then detected by methods well known in the art. Purified GPx6 can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding GPx6 specifically compete with a test compound for binding GPx6. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with GPx6.

In additional embodiments, the nucleotide sequences which encode GPx6 may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

The examples below are provided to illustrate the subject invention and are not included

for the purpose of limiting the invention.

## EXAMPLES

### I. PROSNOT20 cDNA Library Construction

The PROSNOT20 cDNA library was constructed using polyA RNA isolated from nontumorous prostate tissue removed from a 65-year-old Caucasian male during a radical prostatectomy. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated an adenocarcinoma (Gleason grade 2+2) involving the right anterior prostate peripherally. The patient presented with elevated prostate specific antigen (PSA).

The frozen tissue was homogenized and lysed in guanidinium isothiocyanate solution using a Brinkmann Polytron PT-3000 homogenizer (Brinkmann Instruments, Westbury NY). The lysates were centrifuged over a 5.7 M CsCl cushion using a Beckman SW28 rotor in a Beckman L8-70M Ultracentrifuge (Beckman Instruments) for 18 hours at 25,000 rpm at ambient temperature. The RNA was extracted with acid phenol, precipitated using sodium acetate and ethanol, resuspended in RNase-free water, and treated with DNase. RNA extraction and precipitation were repeated as before. Poly(A+) RNA was isolated using the Qiagen Oligotex kit (QIAGEN, Inc., Chatsworth CA).

Poly(A+) RNA was used for cDNA synthesis and construction of the PROSNOT20 cDNA library according to the recommended protocols in the SuperScript plasmid system (Cat. #18248-013, Life Technologies, Gaithersburg MD). cDNAs were fractionated on a Sepharose CL4B column (Cat. #275105-01, Pharmacia Amersham Biotech, Piscataway, NJ) and those cDNAs exceeding 400 bp were ligated into pINCY (Incyte Pharmaceuticals, Inc., Palo Alto CA). The recombinant plasmids were subsequently transformed into DH5a competent cells (Cat. #18258-012, Life Technologies).

### II. Isolation and Sequencing of cDNA Clones

Plasmid DNA was released from the cells and purified using the REAL Prep 96 Plasmid Kit (Catalog #26173, QIAGEN, Inc.). The recommended protocol was employed except for the following changes: 1) the bacteria were cultured in 1 ml of sterile Terrific Broth (Catalog #22711, Life Technologies) with carbenicillin at 25 mg/L and glycerol at 0.4%; 2) after the cultures were incubated for 19 hours, the cells were lysed with 0.3 ml of lysis buffer; and 3) following isopropanol precipitation, the plasmid DNA pellets were resuspended in 0.1 ml of distilled water. The DNA samples were stored at 4°C.

The cDNAs were sequenced by the method of Sanger et al. (1975, *J. Mol. Biol.* 94:441f), using a Hamilton Micro Lab 2200 (Hamilton, Reno NV) in combination with Peltier Thermal Cyclers (PTC200 from MJ Research, Watertown MA) and Applied Biosystems 377 DNA Sequencing Systems, and the reading frame was determined.

### III. Similarity Searching of cDNA Clones and Their Deduced Proteins

The nucleotide sequences and/or amino acid sequences of the Sequence Listing were used to query sequences in the GenBank, SwissProt, BLOCKS, and Pima II databases. These databases, which contain previously identified and annotated sequences, were searched for regions of similarity using BLAST (Basic Local Alignment Search Tool). (See, e.g., Altschul, S.F. (1993) *J. Mol. Evol.* 36:290-300; and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410.)

BLAST produced alignments of both nucleotide and amino acid sequences to determine sequence similarity. Because of the local nature of the alignments, BLAST was especially useful in determining exact matches or in identifying homologs which may be of prokaryotic (bacterial) or eukaryotic (animal, fungal, or plant) origin. Other algorithms could have been used when dealing with primary sequence patterns and secondary structure gap penalties. (See, e.g., Smith, T. et al. (1992) *Protein Eng.* 5:35-51.) The sequences disclosed in this application have lengths of at least 49 nucleotides and have no more than 12% uncalled bases (where N is recorded rather than A, C, G, or T).

The BLAST approach searched for matches between a query sequence and a database sequence. BLAST evaluated the statistical significance of any matches found, and reported only those matches that satisfy the user-selected threshold of significance. In this application, threshold was set at  $10^{-25}$  for nucleotides and  $10^{-8}$  for peptides.

Incyte nucleotide sequences were searched against the GenBank databases for primate (pri), rodent (rod), and other mammalian sequences (mam), and deduced amino acid sequences from the same clones were then searched against GenBank functional protein databases, mammalian (mamp), vertebrate (vrtp), and eukaryote (eukp), for similarity.

Additionally, sequences identified from cDNA libraries may be analyzed to identify those gene sequences encoding conserved protein motifs using an appropriate analysis program, e.g., BLOCKS. BLOCKS is a weighted matrix analysis algorithm based on short amino acid segments, or blocks, compiled from the PROSITE database (Bairoch, A. et al. (1997) *Nucleic Acids Res.* 25:217-221). The BLOCKS algorithm is useful for classifying genes with unknown functions (Henikoff S. and G.J. Henikoff (1991) *Nucleic Acids Res.* 19:6565-6572). Blocks,



which are 3-60 amino acids in length, correspond to the most highly conserved regions of proteins. The BLOCKS algorithm compares a query sequence with a weighted scoring matrix of blocks in the BLOCKS database. Blocks in the BLOCKS database are calibrated against protein sequences with known functions from the SWISS-PROT database to determine the stochastic distribution of matches. Similar databases such as PRINTS, a protein fingerprint database, are also searchable using the BLOCKS algorithm. (Attwood, T.K. et al. (1997) *J. Chem. Inf. Comput. Sci.* 37:417-424.) PRINTS is based on non-redundant sequences obtained from sources such as SWISS-PROT, GenBank, PIR, and NRL-3D.

The BLOCKS algorithm searches for matches between a query sequence and the BLOCKS or PRINTS database and evaluates the statistical significance of any matches found. Matches from a BLOCKS or PRINTS search can be evaluated on two levels, local similarity and global similarity. The degree of local similarity is measured by scores, and the extent of global similarity is measured by score ranking and probability values. A score of 1000 or greater for a BLOCKS match of highest ranking indicates that the match falls within the 0.5 percentile level of false positives when the matched block is calibrated against SWISS-PROT. Likewise, a probability value of less than  $1.0 \times 10^{-3}$  indicates that the match would occur by chance no more than one time in every 1000 searches. Only those matches with a cutoff score of 1000 or greater and a cutoff probability value of  $1.0 \times 10^{-3}$  or less are considered in the functional analyses of the protein sequences in the Sequence Listing.

Nucleic and amino acid sequences of the Sequence Listing may also be analyzed using PFAM. PFAM is a Hidden Markov Model (HMM) based protocol useful in protein family searching. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) *Curr. Opin. Struct. Biol.* 6:361-365.)

The PFAM database contains protein sequences of 527 protein families gathered from publicly available sources, e.g., SWISS-PROT and PROSITE. PFAM searches for well characterized protein domain families using two high-quality alignment routines, seed alignment and full alignment. (See, e.g., Sonnhammer, E.L.L. et al. (1997) *Proteins: Struct. Funct. Genet.* 28:405-420.) The seed alignment utilizes the hmmls program, a program that searches for local matches, and a non-redundant set of the PFAM database. The full alignment utilizes the hmmsf program, a program that searches for multiple fragments in long sequences, e.g., repeats and motifs, and all sequences in the PFAM database. A result or score of 100 "bits" can signify that it is  $2^{100}$ -fold more likely that the sequence is a true match to the model or comparison sequence. Cutoff scores which range from 10 to 50 bits are generally used for individual protein families using the SWISS-PROT sequences as model or comparison sequences.

Two other algorithms, SIGPEPT and TM, both based on the HMM algorithm described above (see, e.g., Eddy, supra; and Sonnhammer, supra), identify potential signal sequences and transmembrane domains, respectively. SIGPEPT was created using protein sequences having signal sequence annotations derived from SWISS-PROT. It contains about 1413 non-redundant signal sequences ranging in length from 14 to 36 amino acid residues. TM was created similarly using transmembrane domain annotations. It contains about 453 non-redundant transmembrane sequences encompassing 1579 transmembrane domain segments. Suitable HMM models were constructed using the above sequences and were refined with known SWISS-PROT signal peptide sequences or transmembrane domain sequences until a high correlation coefficient, a measurement of the correctness of the analysis, was obtained. Using the protein sequences from the SWISS-PROT database as a test set, a cutoff score of 11 bits, as determined above, correlated with 91-94% true-positives and about 4.1% false-positives, yielding a correlation coefficient of about 0.87-0.90 for SIGPEPT. A score of 11 bits for TM will typically give the following results: 75% true positives; 1.72% false positives; and a correlation coefficient of 0.76. Each search evaluates the statistical significance of any matches found and reports only those matches that score at least 11 bits.

#### IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; and Ausubel, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST are used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ database (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar.

The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and

40, although lower scores may identify related molecules.

The results of Northern analysis are reported as a list of libraries in which the transcript encoding GPx6 occurs. Abundance and percent abundance are also reported. Abundance directly reflects the number of times a particular transcript is represented in a cDNA library, and percent abundance is abundance divided by the total number of sequences examined in the cDNA library.

## V. Extension of GPx6 Encoding Polynucleotides

The nucleic acid sequence of Incyte Clone 1817518 was used to design oligonucleotide primers for extending a partial nucleotide sequence to full length. One primer was synthesized to initiate extension of an antisense polynucleotide, and the other was synthesized to initiate extension of a sense polynucleotide. Primers were used to facilitate the extension of the known sequence "outward" generating amplicons containing new unknown nucleotide sequence for the region of interest. The initial primers were designed from the cDNA using OLIGO 4.06 (National Biosciences, Plymouth MN), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries (GIBCO BRL) were used to extend the sequence. If more than one extension is necessary or desired, additional sets of primers are designed to further extend the known region.

High fidelity amplification was obtained by following the instructions for the XL-PCR kit (Perkin Elmer) and thoroughly mixing the enzyme and reaction mix. PCR was performed using the Peltier Thermal Cycler (PTC200; M.J. Research, Watertown MA), beginning with 40 pmol of each primer and the recommended concentrations of all other components of the kit, with the following parameters:

- |         |                                                       |
|---------|-------------------------------------------------------|
| Step 1  | 94° C for 1 min (initial denaturation)                |
| Step 2  | 65° C for 1 min                                       |
| Step 3  | 68° C for 6 min                                       |
| Step 4  | 94° C for 15 sec                                      |
| Step 5  | 65° C for 1 min                                       |
| Step 6  | 68° C for 7 min                                       |
| Step 7  | Repeat steps 4 through 6 for an additional 15 cycles  |
| Step 8  | 94° C for 15 sec                                      |
| Step 9  | 65° C for 1 min                                       |
| Step 10 | 68° C for 7:15 min                                    |
| Step 11 | Repeat steps 8 through 10 for an additional 12 cycles |

Step 12	72° C for 8 min
Step 13	4° C (and holding)

A 5  $\mu$ l to 10  $\mu$ l aliquot of the reaction mixture was analyzed by electrophoresis on a low concentration (about 0.6% to 0.8%) agarose mini-gel to determine which reactions were successful in extending the sequence. Bands thought to contain the largest products were excised from the gel, purified using QIAQUICK (QIAGEN Inc.), and trimmed of overhangs using Klenow enzyme to facilitate religation and cloning.

After ethanol precipitation, the products were redissolved in 13  $\mu$ l of ligation buffer, 1  $\mu$ l T4-DNA ligase (15 units) and 1  $\mu$ l T4 polynucleotide kinase were added, and the mixture was incubated at room temperature for 2 to 3 hours, or overnight at 16° C. Competent *E. coli* cells (in 40  $\mu$ l of appropriate media) were transformed with 3  $\mu$ l of ligation mixture and cultured in 80  $\mu$ l of SOC medium. (See, e.g., Sambrook, supra, Appendix A, p. 2.) After incubation for one hour at 37°C, the *E. coli* mixture was plated on Luria Bertani (LB) agar (See, e.g., Sambrook, supra, Appendix A, p. 1) containing carbenicillin (2x carb). The following day, several colonies were randomly picked from each plate and cultured in 150  $\mu$ l of liquid LB/2x carb medium placed in an individual well of an appropriate commercially-available sterile 96-well microtiter plate. The following day, 5  $\mu$ l of each overnight culture was transferred into a non-sterile 96-well plate and, after dilution 1:10 with water, 5  $\mu$ l from each sample was transferred into a PCR array.

For PCR amplification, 18  $\mu$ l of concentrated PCR reaction mix (3.3x) containing 4 units of rTth DNA polymerase, a vector primer, and one or both of the gene specific primers used for the extension reaction were added to each well. Amplification was performed using the following conditions:

Step 1	94° C for 60 sec
Step 2	94° C for 20 sec
Step 3	55° C for 30 sec
Step 4	72° C for 90 sec
Step 5	Repeat steps 2 through 4 for an additional 29 cycles
Step 6	72° C for 180 sec
Step 7	4° C (and holding)

Aliquots of the PCR reactions were run on agarose gels together with molecular weight markers. The sizes of the PCR products were compared to the original partial cDNAs, and appropriate clones were selected, ligated into plasmid, and sequenced.

In like manner, the nucleotide sequence of SEQ ID NO:2 is used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for 5' extension, and an appropriate genomic library.

## VI. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:2 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250  $\mu$ Ci of [ $\gamma$ - $^{32}$ P] adenosine triphosphate (Amersham, Chicago IL), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Pharmacia & Upjohn, Kalamazoo MI). An aliquot containing  $10^7$  counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN, Boston MA).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under increasingly stringent conditions up to 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. After XOMAT AR film (Kodak, Rochester NY) is exposed to the blots to film for several hours, hybridization patterns are compared visually.

## VII. Microarrays

A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Baldeschweiler, supra.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE. Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or

selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470; and Shalon, D. et al. (1996) *Genome Res.* 6:639-645.)

Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

### VIII. Complementary Polynucleotides

Sequences complementary to the GPx6-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring GPx6. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software and the coding sequence of GPx6. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the GPx6-encoding transcript.

### IX. Expression of GPx6

Expression and purification of GPx6 is achieved using bacterial or virus-based expression systems. For expression of GPx6 in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express GPx6 upon induction with isopropyl beta-D-thiogalactopyranoside (IPTG). Expression of GPx6 in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant *Autographica californica* nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding GPx6 by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect *Spodoptera frugiperda* (Sf9) insect cells in most cases,

or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, GPx6 is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from *Schistosoma japonicum*, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Pharmacia, Piscataway, NJ). Following purification, the GST moiety can be proteolytically cleaved from GPx6 at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak, Rochester NY). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN Inc, Chatsworth CA). Methods for protein expression and purification are discussed in Ausubel, F.M. et al. (1995 and periodic supplements) *Current Protocols in Molecular Biology*, John Wiley & Sons, New York NY, ch. 10, 16. Purified GPx6 obtained by these methods can be used directly in the following activity assay.

## X. Demonstration of GPx6 Activity

GPx6 activity is measured spectrophotometrically by a procedure described by Maiorino et al. (1990; *supra*). Reaction substrate is prepared by oxygenation of soybean phosphatidylcholine using soybean lipoxidase in the presence of bile salts. After 50% of the oxygen has been consumed, the phospholipids are separated from the bile salts by column chromatography. Cell homogenates are assayed for GPx6 activity by mixing homogenate with peroxide-free Triton X-100, an excess amount of glutathione reductase, and saturating amounts of NADPH in a buffer containing reduced glutathione. Glutathione oxidation is recorded at 340 nm at baseline (before substrate addition) and following substrate addition until reaction rate returns to baseline. One unit of activity catalyzes the reduction of 1  $\mu\text{mol}$  glutathione/min at 37 °C.

## XI. Functional Assays

GPx6 function is assessed by expressing the sequences encoding GPx6 at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian

expression vector containing a strong promoter that drives high levels of cDNA expression.

Vectors of choice include pCMV SPORT (Life Technologies, Gaithersburg MD) and pCR 3.1 (Invitrogen, Carlsbad, CA, both of which contain the cytomegalovirus promoter. 5-10  $\mu$ g of recombinant vector are transiently transfected into a human cell line, preferably of endothelial or hematopoietic origin, using either liposome formulations or electroporation. 1-2  $\mu$ g of an additional plasmid containing sequences encoding a marker protein are co-transfected.

Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP) (Clontech, Palo Alto CA), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP, and to evaluate properties, for example, their apoptotic state. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of GPx6 on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding GPx6 and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding GPx6 and other genes of interest can be analyzed by Northern analysis or microarray techniques.

## **XII. Production of GPx6 Specific Antibodies**

GPx6 substantially purified using polyacrylamide gel electrophoresis (PAGE) (see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.



Alternatively, the GPx6 amino acid sequence is analyzed using LASERGENE software (DNASTAR Inc.) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel supra, ch. 11.)

Typically, oligopeptides 15 residues in length are synthesized using an Applied Biosystems Peptide Synthesizer Model 431A using fmoc-chemistry and coupled to KLH (Sigma, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide activity by, for example, binding the peptide to plastic, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

### XIII. Purification of Naturally Occurring GPx6 Using Specific Antibodies

Naturally occurring or recombinant GPx6 is substantially purified by immunoaffinity chromatography using antibodies specific for GPx6. An immunoaffinity column is constructed by covalently coupling anti-GPx6 antibody to an activated chromatographic resin, such as CNBr-activated Sepharose (Pharmacia & Upjohn). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing GPx6 are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of GPx6 (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/GPx6 binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and GPx6 is collected.

### XIV. Identification of Molecules Which Interact with GPx6

GPx6, or biologically active fragments thereof, are labeled with <sup>125</sup>I Bolton-Hunter reagent. (See, e.g., Bolton, A.E. and W.M. Hunter (1973) *Biochem. J.* 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled GPx6, washed, and any wells with labeled GPx6 complex are assayed. Data obtained using different concentrations of GPx6 are used to calculate values for the number, affinity, and association of GPx6 with the candidate molecules.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
22

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: HILLMAN, JENNIFER L.  
CORLEY, NEIL C.  
PATTERSON, CHANDRA
- (ii) TITLE OF INVENTION: HUMAN GLUTATHIONE PEROXIDASE-6
- (iii) NUMBER OF SEQUENCES: 3
- (iv) CORRESPONDENCE ADDRESS:  
(A) ADDRESSEE: Incyte Pharmaceuticals, Inc.  
(B) STREET: 3174 Porter Drive  
(C) CITY: Palo Alto  
(D) STATE: CA  
(E) COUNTRY: USA  
(F) ZIP: 94304
- (v) COMPUTER READABLE FORM:  
(A) MEDIUM TYPE: Diskette  
(B) COMPUTER: IBM Compatible  
(C) OPERATING SYSTEM: Windows  
(D) SOFTWARE: FastSEQ for Windows Version 2.0b
- (vi) CURRENT APPLICATION DATA:  
(A) APPLICATION NUMBER: TO BE ASSIGNED  
(B) FILING DATE: HEREWITH  
(C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:  
(A) APPLICATION NUMBER:  
(B) FILING DATE:
- (viii) ATTORNEY/AGENT INFORMATION:  
(A) NAME: Cerrone, Michael C  
(B) REGISTRATION NUMBER: 39,132  
(C) REFERENCE/DOCKET NUMBER: PF-0530 US
- (ix) TELECOMMUNICATION INFORMATION:  
(A) TELEPHONE: 650-855-0555  
(B) TELEFAX: 650-855-0572  
(C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 187 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear
- (vii) IMMEDIATE SOURCE:  
(A) LIBRARY: PROSNOT20  
(B) CLONE: 1817518

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

Met Val Ala Ala Thr Val Ala Ala Ala Trp Leu Leu Leu Trp Ala Ala
 1          5          10
Ala Cys Ala Gln Gln Glu Gln Asp Phe Tyr Asp Phe Lys Ala Val Asn
 20          25          30
Ile Arg Gly Lys Leu Val Ser Leu Glu Lys Tyr Arg Gly Ser Val Ser
 35          40          45
Leu Val Val Asn Val Ala Ser Glu Cys Gly Phe Thr Asp Gln His Tyr
 50          55          60
Arg Ala Leu Gln Gln Leu Gln Arg Asp Leu Gly Pro His His Phe Asn
 65          70          75
Val Leu Ala Phe Pro Cys Asn Gln Phe Gly Gln Gln Glu Pro Asp Ser
 85          90          95
Asn Lys Glu Ile Glu Ser Phe Ala Cys Arg Thr Tyr Ser Val Ser Phe
100          105          110
Pro Met Phe Ser Lys Ile Ala Val Thr Gly Thr Gly Ala His Pro Ala
115          120          125
Phe Lys Tyr Leu Ala Gln Thr Ser Gly Lys Glu Pro Thr Trp Asn Phe
130          135          140
Trp Lys Tyr Leu Val Ala Pro Asp Gly Lys Val Val Gly Ala Trp Asp
145          150          155
Pro Thr Val Ser Val Glu Glu Val Arg Leu Gln Ile Thr Ala Leu Val
165          170          175
Arg Lys Leu Ile Leu Leu Lys Arg Glu Asp Leu
180          185

```

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1072 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

## (vii) IMMEDIATE SOURCE:

- (A) LIBRARY: PROSNOT20  
 (B) CLONE: 1817518

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

```

GACGCGGCCA CCTCCGGAAC AAGCCATGGT GCGGCGGACG GTGGCAGCGG CGTGGCTGCT 60
CCTGTGGGCT GCGGCCTGCG CGCAGCAGGA GCAGGACTTC TACGACTTCA AGGCGGTCAA 120
CATCCCGGGC AAACGTGGTGT CGCTGGAGAA GTACCGCGGA TCGGTGTCCC TGGTGGTGAA 180
TGTTGCGCAGC GAGCTGCGGCT TCACAGACCA GCCTACCGA GCCCTGCAGC AGCTGCAGCG 240
AGACCTGGGC CCCCACCACT TCAACGTGCT CGCCTTCCCC TGCAACCACT TTGGCCAACA 300
GGAGCCTGAC AGCAACAAGG AGATTGAGAG CTTTGCCTGC CGCACCTACA GTGTCTCAAT 360
CCCCATGTTT AGCAAGATTG CAGTCACCGG TACTGGTGCC CATCTGCTCT TCAAGTACCT 420
GGCCAGACTC TCTGGAAGAG AGCCACCTCG GAACCTCTGG AAGTACCTAG TAGCCCCAGA 480
TGGAAAGGTT GTAGGGGCTT GGGACCCCAAC TGTGTCACTG GAGGAGGTCA GACTCCAGAT 540
CACAGCGCTC CTGAGGAAGC TCATCTTACT GAAGCGAGAA GACTTATAAC CACCGCGTCT 600
CCTCCTCCAC CACCTCATCC CGCCACCTCG TGTGGGGCTG ACCAATGCCA ACTCAAAATG 660
TGCTTCAAAG GGAGAGACCC ACTGACTCTC CTTCCTTAC TCTTATGCCA TTGGTCCCAT 720
CATTCTTTGT GGGGAAAAAT TCTAGTATTT TGATATTATT AATCTTACAG CAACAAATAG 780
GAACTCTGGG CCAATGAGAG CTCTTGACCA GTGAATCACC AGCCGATACG AACGCTTCGC 840
CAACAAAAAT GTGTGCAAAA TAGAAGTATA TCAAGCAATA ATCTCCCAAC CAAGGCTTCT 900
GTAAACTGGG ACCAATGATT ACCTCATAGG GCTGTTGTGA GGATTAGGAT GAAATACCTG 960
TGAAAGTGCC TAGGCAGTGC CAGCCAAATA GGAGGCATTC AATGAACATT TTTTGCATAT 1020
AAACCAAAAA ATAACCTGTT ATCAATAAAA ACTTGCAATC AACATGAATT TC 1072

```

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 170 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY: GENBANK
- (B) CLONE: 544432

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Met	Cys	Ala	Ser	Arg	Asp	Asp	Trp	Arg	Cys	Ala	Arg	Ser	Met	His	Glu
1				5					10					15	
Phe	Ser	Ala	Lys	Asp	Ile	Asp	Gly	His	Met	Val	Asn	Leu	Asp	Lys	Tyr
			20					25					30		
Arg	Gly	Phe	Val	Cys	Ile	Val	Thr	Asn	Val	Ala	Ser	Gln	Cys	Gly	Lys
			35					40					45		
Thr	Glu	Val	Asn	Tyr	Thr	Gln	Leu	Val	Asp	Leu	His	Ala	Arg	Tyr	Ala
			50					55				60			
Glu	Cys	Gly	Leu	Arg	Ile	Leu	Ala	Phe	Pro	Cys	Asn	Gln	Phe	Gly	Lys
			65			70			75					80	
Gln	Glu	Pro	Gly	Ser	Asn	Glu	Glu	Ile	Lys	Glu	Phe	Ala	Ala	Gly	Tyr
				85					90					95	
Asn	Val	Lys	Phe	Asp	Met	Phe	Ser	Lys	Ile	Cys	Val	Asn	Gly	Asp	Asp
			100					105					110		
Ala	His	Pro	Leu	Trp	Lys	Trp	Met	Lys	Ile	Gln	Pro	Lys	Gly	Lys	Gly
			115				120					125			
Ile	Leu	Gly	Asn	Ala	Ile	Lys	Trp	Asn	Phe	Thr	Lys	Phe	Leu	Ile	Asp
			130			135					140				
Lys	Asn	Gly	Cys	Val	Val	Lys	Arg	Tyr	Gly	Pro	Met	Glu	Glu	Pro	Leu
			145			150				155				160	
Val	Ile	Glu	Lys	Asp	Leu	Pro	His	Tyr	Phe						
				165					170						